

# Siam weed (*Chromolaena odorata* L.) for phytoremediation of artisanal gold mine tailings

*by* A Hamzah

---

**Submission date:** 06-Jun-2020 05:01AM (UTC+0300)

**Submission ID:** 1338702659

**File name:** 2.\_JOURNAL,\_2012.\_JTROPAG\_AHZ.docx (63.86K)

**Word count:** 1480

**Character count:** 7393



## Short communication

# Siam weed (*Chromolaena odorata* L.) for phytoremediation of artisanal gold mine tailings

A. Hamzah<sup>1</sup>,

## Abstract

The potential of *Chromolaena odorata* to remove Hg and Pb from tailings from cyanidation processing of artisanal gold mining at Lombok, Indonesia was investigated in a glasshouse experiment. The tailings were amended with farmyard manure (FYM) and biochar added with 5 and 10 g kg<sup>-1</sup> FeSO<sub>4</sub> before growing *C. odorata* plants. *Chromolaena odorata* grew well on amended tailings, but it did not grow on untreated tailings. Application of organic amendments and FeSO<sub>4</sub> decreased tailing pH from 9.2 up to 6.6 (e.g., FYM+10 g FeSO<sub>4</sub> kg<sup>-1</sup> soil) and increased its organic C and N, P, K levels. Total absorption of Hg in *C. odorata* plants ranged from 5.8 to 8.7 mg kg<sup>-1</sup> and Pb from 55 to 61 mg kg<sup>-1</sup> in the amended substrates.

**Keywords:** Organic amendments, Biochar, Artisanal mining

Small-scale gold mining called artisanal mining is increasing in Indonesia and is estimated to reach 713 spots in Sumatra, Java, Kalimantan, Sulawesi and East Nusa Tenggara. Although mining activities have been capable of contributing to employment and improve societal welfare, artisanal mining has led to environmental problems due to heavy metal contamination. Artisanal mining in Indonesia mostly use the cheap and easy mercury amalgamation technique for gold recovery, which results in mercury contamination of the environment. Krisnayanti et al. (2012) reported as high as 3000 mg kg<sup>-1</sup> Hg in amalgamation tailings and 1900 mg kg<sup>-1</sup> in cyanide tailings in Lombok, Indonesia. Phytoremediation has been widely recommended for removing heavy metals from the environment. The plants used for reclaiming contaminated sites are called “hyper accumulators”, which absorb and accumulate high amount of metal without experi-

encing phytotoxicity and are therefore capable of growing in unfavorable medium and produce high biomass yields. Hartley et al. (2009) observed very low organic C content and very high pH in the tailing of artisanal gold mining at Sekotong, Lombok, Indonesia, which necessitates addition of soil amendments such as farmyard manure (FYM) or more resistant organic sources such as biochar in conjunction with sulfur containing materials, such as elemental sulfur, Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>, or FeSO<sub>4</sub>. This study examined the potential of *Chromolaena odorata* plants to remove heavy metals, especially mercury (Hg) and lead (Pb) from tailing of artisanal gold mining, following additions of various soil amendments.

The glasshouse experiment was carried out at the Tribhuwana Tunggal University, Malang, Indonesia. Tailing of cyanidation gold processing

were collected from artisanal gold mining at West Lombok, Indonesia. These tailings were air-dried, crushed, and passed through a 2.0 mm diameter sieve. The treatments included: (1) FYM + 5 g  $\text{FeSO}_4$  kg<sup>-1</sup> soil, (2) FYM + 10 g  $\text{FeSO}_4$  kg<sup>-1</sup> soil, (3) biochar + 5 g  $\text{FeSO}_4$  kg<sup>-1</sup> soil, (4) biochar + 10 g  $\text{FeSO}_4$  kg<sup>-1</sup> soil, and (5) no amendment as control, arranged in a completely randomized design with three replications. FYM was obtained from a local farmer at Malang, Indonesia and the biochar was prepared by pyrolysis of FYM. Some characteristics of tailings, FYM, and biochar used in this experiment are presented in Table 1.

Ten kg of crushed air-dried tailing (water content of about 14%) was put in a plastic pot (15 kg capacity), watered to about field capacity, incubated for 30 days, and planted with *C. odorata*. Ten gram N, P, and K fertilizers from urea (46% N), super phosphate 36 (36%  $\text{P}_2\text{O}_5$ ), and potassium chloride (50%  $\text{K}_2\text{O}$ ) were applied at planting time, and the plants were grown in a glasshouse for four months. During the experiment, water content was maintained at about field capacity by daily watering. Soil samples were collected 30 days after incubation for analysis of pH, organic C, total mercury (Hg), total lead (Pb). Plant height, leaf number, dry biomass, and the content of Hg and Pb were determined at harvest. The plant samples (leaves, stems, and roots) were washed with tap water, rinsed with deionized water, and oven-dried at 60–70°C for 72 h. The samples were ground in an electric mill and homogenized by thorough mixing.

Total Hg and Pb in soils and plants were extracted using  $\text{HNO}_3$  and  $\text{HClO}_4$  and soluble Hg and Pb using  $\text{HCl}$ ,  $\text{HNO}_3$ , and  $\text{K}_2\text{Cr}_2\text{O}_7$ , and the concentration of Hg and Pb determined by Atomic Absorption Spectrometry (AAS). Soil pH was measured in 1:2.5 ratio of soil and de-ionized water. The Walkley and Black wet oxidation method determined soil organic C. The data were analyzed using ANOVA in Minitab Program ( $p=0.05$ ).

Solubility of Hg and Pb were not significantly influenced by soil amendments (Table 2). Theoretically, a decrease in tailing pH would increase the solubility of heavy metals, but it seems that the pH changes in this study (up to 6.6) did not influence metal solubility. Application of amendments significantly influenced the tailing pH, organic C, available P, exchangeable K, and N content (Table 2). Both FYM and biochar application increased organic C in the tailings. Tailing pH decreased with application of  $\text{FeSO}_4$ , and pH change was higher as the  $\text{FeSO}_4$  rate increased. The pH changes in FYM applied tailing was higher compared to that in biochar treated tailing. This might be due to a lower FYM pH (Table 1). The results in Table 2 further show that the content of P and K increased as pH approached neutrality, which is normal. Availability of P in alkaline condition is generally low because of P fixation by calcium.

*Chromolaena odorata* did not survive on untreated tailings. Even on amended tailings, it exhibited ab-

Table 1. Tailing, farmyard manure (FYM), and biochar characteristics from West Lombok, Indonesia.

| Properties                   | Tailing | Farmyard manure | Biochar |
|------------------------------|---------|-----------------|---------|
| pH ( $\text{H}_2\text{O}$ )  | 9.1     | 6.8             | 8.2     |
| Organic C (%)                | 0.21    | 13.48           | 25.77   |
| Total N (%)                  | 0.13    | 1.14            | 0.52    |
| K (cmol kg <sup>-1</sup> )   | 0.11    | 1.00            | 1.23    |
| CEC (cmol kg <sup>-1</sup> ) | 13.23   | –               | 39.62   |
| Hg (mg kg <sup>-1</sup> )    | 247.24  | –               | –       |
| Pb (mg kg <sup>-1</sup> )    | 134.26  | –               | –       |

CEC: Cation exchange capacity

Table 2. Some characteristics of tailings from West Lombok, Indonesia treated with amendments 30 days after incubation.

| Treatment  | pH    | C<br>(mg kg <sup>-1</sup> ) | Total N<br>(mg kg <sup>-1</sup> ) | Total P<br>(mg kg <sup>-1</sup> ) | Exchangeable K<br>(cmol kg <sup>-1</sup> ) | Soluble Hg<br>(mg kg <sup>-1</sup> ) | Soluble Pb<br>(mg kg <sup>-1</sup> ) |
|--|-------|-----------------------------|-----------------------------------|-----------------------------------|--|--------------------------------------|--------------------------------------|
| FYM + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil      | 7.3ab | 0.29b                       | 0.02b                             | 92.4bc                            | 4.5c                                       | 0.3a                                 | 9.3a                                 |
| FYM + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil     | 6.6a  | 0.25b                       | 0.02b                             | 119.9c                            | 5.6c                                       | 0.4a                                 | 10.9a                                |
| Biochar + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil  | 7.4b  | 0.36c                       | 0.20b                             | 60.7b                             | 1.9b                                       | 0.2a                                 | 10.2a                                |
| Biochar + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil | 7.1ab | 0.36c                       | 0.02b                             | 89.1b                             | 2.5b                                       | 0.1a                                 | 10.3a                                |
| Control  | 9.2c  | 0.02a                       | 0.008a                            | 12.4a                             | 0.1a                                       | 0 0                                  |                                      |

Means followed by the same letters, in same column was not significantly different (p=0.05)

normal growth characteristics (curly and chlorotic leaves, with brown spots). As can be seen from Table 3, plants grew well in the FYM treated tailing than biochar treated tailing, presumably due to the better fertility status of tailing treated with FYM. Growth of *C. odorata* improved as the rate of FeSO<sub>4</sub> increased, this was reasonable, because the pH approached neutrality (Table 2). Hg and Pb uptake were significantly influenced by organic amendments (Table 4). In general, Hg and Pb absorption of *C. odorata* in FYM treated substrate was higher compared to that of biochar treated tailing. Hg and

Pb absorption also increased with increasing rate of FeSO<sub>4</sub> in the biochar treatment owing to pH changes (Table 2). It is widely known that the solubility of most heavy metals increases as soil pH decreases. In addition, the higher CEC of biochar would make the affinity of biochar to any chemical compound higher compared to FYM, which in turn decreased the availability of metals for plant growth. More than 50% of the absorbed metals accumulated in the foliage—with only a relatively small proportion in stems and roots.

Table 3. Growth characteristics of 120-day-old *Chromolaena odorata* on tailings from West Lombok, Indonesia treated with amendments.

| Treatments   | Plant height (cm) | Leaf number/plant | Dry biomass g/plant |
|--|-------------------|-------------------|---------------------|
| FYM + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil      | 112.8b            | 112b              | 97.7b               |
| FYM + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil     | 119.4b            | 129b              | 128.8c              |
| Biochar + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil  | 84.6a             | 90a               | 62.0a               |
| Biochar + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil | 87.9a             | 91a               | 68.0a               |

Means followed by the same letters, in same column was not significantly different (p=0.05)

Table 4. Effect of FYM, biochar and FeSO<sub>4</sub> application on the uptake of Hg and Pb by 120-day-old *Chromolaena odorata* plants grown on tailings from West Lombok, Indonesia treated with amendments.

| Treatments   | Mercury (Hg)                    |               |             |              | Lead (Pb)                       |               |             |              |
|--|---------------------------------|---------------|-------------|--------------|---------------------------------|---------------|-------------|--------------|
|  | Total<br>(mg kg <sup>-1</sup> ) | Leaves<br>(%) | Stem<br>(%) | Roots<br>(%) | Total<br>(mg kg <sup>-1</sup> ) | Leaves<br>(%) | Stem<br>(%) | Roots<br>(%) |
| FYM + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil      | 8.9b                            | 53.5a         | 22.6a       | 23.9a        | 57.0ab                          | 55.6a         | 20.2a       | 24.2a        |
| FYM + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil     | 8.7b                            | 52.1a         | 24.6a       | 23.3a        | 58.2ab                          | 56.2a         | 15.6a       | 28.2a        |
| Biochar + 5 g FeSO <sub>4</sub> kg <sup>-1</sup> soil  | 5.8a                            | 57.0a         | 20.6a       | 22.4a        | 55.0a                           | 55.3a         | 15.2a       | 29.5a        |
| Biochar + 10 g FeSO <sub>4</sub> kg <sup>-1</sup> soil | 8.0b                            | 52.5a         | 21.5a       | 26.0a        | 61.0b                           | 57.3a         | 18.6a       | 24.1a        |

Means followed by the same letters, in same column was not significantly different (p=0.05)

Application of organic amendments in conjunction with  $\text{FeSO}_4$  decreased tailing pH from 9.2 up to 6.6 (FYM+10 g  $\text{FeSO}_4 \text{ kg}^{-1}$  soil), besides increasing organic C and N, P, and K levels. *Chromolaena odorata* has a good potential for removal of Hg and Pb from artisanal gold mining, because of its ability to accumulate high concentrations of these metals without experiencing toxicity. Total absorption of Hg varied from 5.8 (FYM+5 g  $\text{FeSO}_4 \text{ kg}^{-1}$  soil) to 8.7  $\text{mg kg}^{-1}$  (FYM+10 g  $\text{FeSO}_4 \text{ kg}^{-1}$  soil), and Pb from 55.0 (biochar+5 g  $\text{FeSO}_4 \text{ kg}^{-1}$  soil) to 61.0  $\text{mg kg}^{-1}$  (biochar+10 g  $\text{FeSO}_4 \text{ kg}^{-1}$  soil).



# Siam weed (*Chromolaena odorata* L.) for phytoremediation of artisanal gold mine tailings

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- 1

Chinh T. Tran, Thoa T. Le, Lim T. Duong, Stefan Dultz, Minh N. Nguyen. "A comparative study of arsenic in rice in lowland and terraced paddies in the Red River basin, Vietnam", *Land Degradation & Development*, 2020  
Publication

2%
- 2

[www.koko.gov.my](http://www.koko.gov.my)  
Internet Source

1%
- 3

Trajče Stafilov, Robert Šajn, Zlatko Pančevski, Blažo Boev, Marina V. Frontasyeva, Lyudmila P. Strelkova. "Heavy metal contamination of topsoils around a lead and zinc smelter in the Republic of Macedonia", *Journal of Hazardous Materials*, 2010  
Publication

1%
- 4

Louisa J. Esdaile, Justin M. Chalker. "The Mercury Problem in Artisanal and Small-Scale Gold Mining", *Chemistry - A European Journal*, 2018  
Publication

1%
- 5

[arid.lzu.edu.cn](http://arid.lzu.edu.cn)  
Internet Source

|    |   |     |
|----|---|-----|
|    |   | 1 % |
| 6  | <a href="http://mro.massey.ac.nz">mro.massey.ac.nz</a><br>Internet Source   | 1 % |
| 7  | Liao, M.. "Effect of heavy metals on substrate utilization pattern, biomass, and activity of microbial communities in a reclaimed mining wasteland of red soil area", Ecotoxicology and Environmental Safety, 200702<br>Publication | 1 % |
| 8  | <a href="http://www.virtualacademia.com">www.virtualacademia.com</a><br>Internet Source   | 1 % |
| 9  | Submitted to Universitas Brawijaya<br>Student Paper   | 1 % |
| 10 | <a href="http://www.aensiweb.net">www.aensiweb.net</a><br>Internet Source   | 1 % |
| 11 | <a href="http://jtropag.in">jtropag.in</a><br>Internet Source   | 1 % |
| 12 | <a href="http://ijoeear.com">ijoeear.com</a><br>Internet Source   | 1 % |
| 13 | Fernanda Pereira, Inoka Amarakoon, Francis Zvomuya, Nicholson Jeke. "Kinetics and Thermodynamics of Phosphorus Sorption on Goethites: Effects of Biochar Application", Canadian Journal of Soil Science, 2017<br>Publication        | 1 % |

Ali El-Naggar, Sang Soo Lee, Jörg Rinklebe,

|    |  |      |
|----|--|------|
| 14 | Muhammad Farooq et al. "Biochar application to low fertility soils: A review of current status, and future prospects", Geoderma, 2019<br>Publication   | 1 %  |
| 15 | <a href="http://www.tandfonline.com">www.tandfonline.com</a><br>Internet Source  | 1 %  |
| 16 | R. L. Reed <sup>1</sup> , M. A. Sanderson, V. G. Allen, R. E. Zartman. "Cadmium application and pH effects on growth and cadmium accumulation in switchgrass", Communications in Soil Science and Plant Analysis, 2006<br>Publication                              | 1 %  |
| 17 | Moreno, A.M.. "Factors influencing lead sorption-desorption at variable added metal concentrations in Rhodoxeralfs", Chemosphere, 200607<br>Publication  | 1 %  |
| 18 | Cassie R. Rodrigues, Bernard F. Rodrigues. "Influence of organic amendments and arbuscular mycorrhizal inoculation on plant growth and nutrient uptake in iron ore mine spoils", International Journal of Mining, Reclamation and Environment, 2014<br>Publication | <1 % |
| 19 | Submitted to University of Greenwich<br>Student Paper  | <1 % |



Exclude quotes      On

Exclude bibliography      On

Exclude matches      Off

# Siam weed (Chromolaena odorata L.) for phytoremediation of artisanal gold mine tailings

## GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4